



**QUEEN'S
UNIVERSITY
BELFAST**

Arsenic in Bangladeshi soils related to physiographic region, paddy management, and mirco- and macro-elemental status

Chowdhury, M. T. A., Deacon, C. M., Jones, G. D., Imamul Huq, S. M., Williams, P. N., Manzurul Hoque, A. F. M., Winkel, L. H. E., Price, A. H., Norton, G. J., & Meharg, A. A. (2017). Arsenic in Bangladeshi soils related to physiographic region, paddy management, and mirco- and macro-elemental status. *The Science of the total environment*, 590-591, 406-415.

Published in:

The Science of the total environment

Document Version:

Peer reviewed version

Queen's University Belfast - Research Portal:

[Link to publication record in Queen's University Belfast Research Portal](#)

Publisher rights

© 2016 Elsevier B.V. All rights reserved. This manuscript version is made available under the CC-BY-NC-ND 4.0 license <http://creativecommons.org/licenses/by-nc-nd/4.0/>, which permits distribution and reproduction for non-commercial purposes, provided the author and source are cited.

General rights

Copyright for the publications made accessible via the Queen's University Belfast Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The Research Portal is Queen's institutional repository that provides access to Queen's research output. Every effort has been made to ensure that content in the Research Portal does not infringe any person's rights, or applicable UK laws. If you discover content in the Research Portal that you believe breaches copyright or violates any law, please contact openaccess@qub.ac.uk.

**Arsenic in Bangladeshi soils related to physiographic region, paddy
management, and mirco- and macro- elemental status**

M. Tanvir A. Chowdhury^{1, 2}, Claire M. Deacon¹, Gerrad D. Jones³, S.M. Imamul Huq², Paul N. Williams⁴, A.F.M. Manzurul Hoque⁵, Lenny H.E. Winkel⁶, Adam H. Price¹, Gareth J. Norton^{1*}, and Andrew A. Meharg^{4*}

¹Institute of Biological and Environmental Sciences, University of Aberdeen, Cruickshank Building, St. Machar Drive, Aberdeen, AB24 3UU, U.K.

²Department of Soil, Water and Environment, University of Dhaka, Dhaka-1000, Bangladesh

³Eawag: Swiss Federal Institute of Aquatic Science and Technology, Ueberlandstrasse 133, P.O. Box 611, CH-8600 Duebendorf, Switzerland

⁴Institute for Global Food Security, Queen's University Belfast, David Keir Building, Malone Road, Belfast BT9 5BN, Northern Ireland, U.K.

⁵Soil Resource Development Institute (SRDI), Khamar Bari Road, Dhaka 1215, Bangladesh

⁶Swiss Federal Institute of Technology (ETH), Institute of Biogeochemistry and Pollutant Dynamics, ETH Zurich, CH-8092 Zurich, Switzerland

Corresponding authors

*** Andrew A. Meharg**

Tel: +44 (0) 28 9097 5413; fax: +44 (0) 28 9097 6513; e-mail: aa.meharg@qub.ac.uk

*** Gareth J. Norton**

Tel: +44 (0) 1224 272409; fax: +44 (0) 1224 272396; e-mail: g.norton@abdn.ac.uk

Abstract

While the impact of arsenic in irrigated agriculture has become a major environmental concern in Bangladesh, to date there is still a limited understanding of arsenic in Bangladeshi paddy soils at a landscape level. A soil survey was conducted across ten different physiographic regions of Bangladesh, which encompassed six types of geomorphology (Bil, Brahmaputra floodplain, Ganges floodplain, Meghna floodplain, Karatoya-Bangali floodplain and Pleistocene terrace). A total of 1209 paddy soils and 235 matched non-paddy soils were collected. The source of irrigation water (groundwater and surface water) was also recorded. The concentrations of arsenic and sixteen other elements were determined in the soil samples. The concentration of arsenic was higher in paddy soils compared to non-paddy soils, with soils irrigated with groundwater being higher in arsenic than those irrigated with surface water. There was a clear difference between the Holocene floodplains and the Pleistocene terraces, with Holocene floodplain soils being higher in arsenic and other elements. The results suggest that arsenic is most likely associated with less well weathered/ leached soils, suggesting it is either due to the geological newness of Holocene sediments or differences between the sources of sediments, which gives rise to the arsenic problems in Bangladeshi soils.

1. Introduction

Rice is elevated in inorganic arsenic compared to all other dietary staples (Meharg et al., 2009). Flooding of soils, as in paddy cultivation, leads to the mobilization of natural and anthropogenic inorganic arsenic stored in iron oxyhydroxide phases, caused by both the reduction of arsenic and iron under negative soil redox potentials (Meharg and Zhao, 2012). Paddy soils are managed through tilling, fertilization, and surface water and groundwater irrigation, with the latter often elevated in inorganic arsenic throughout large areas of Bangladesh (Huq et al., 2003; Meharg and Rahman, 2003; Roberts et al., 2007; Lu et al., 2009). Furthermore, arsenic can undergo a number of processes within paddy soils that lead to its subsequent loss such as partitioning to monsoonal floodwaters (Dittmar et al., 2007; Saha and Ali, 2007; Dittmar et al., 2010; Roberts et al., 2010), leaching to sub-surfaces (McLaren et al., 2006; Khan et al., 2009; Heikens et al., 2007), and biovolatilization to arsines (Mestrot et al., 2011). Thus, the arsenic loading of any particular paddy soil will be due to geological origin and the subsequent weathering of constituent minerals, and the agronomic management of that sediment (Lu et al., 2009).

Bangladesh has three major geomorphological units (Brammer, 1996; Huq and Shoaib, 2013). These are hill, terrace, and floodplain areas. The hills occupy twelve percent of the country's land area. The uplifted terrace areas are of Pleistocene age and occupy eight percent of the country. The floodplains are of Holocene age and occupy eighty percent of the country. The Holocene floodplains include the piedmont plains, river floodplains, tidal floodplains, and estuarine floodplains. These geomorphological units are related to the

parent geological formations, however, they are also characterized by land topography and age of the soil formation through sediment deposition over time (Brammer, 1996).

To understand and characterise the physiography of the geomorphological areas, Bangladesh is divided into twenty main physiographic regions (FAO/UNDP, 1988). This physiographic classification was based on the parent material in which individual soil types were formed and the landscape on which the soils were developed (FAO/UNDP, 1988). Therefore, the physiographic regions have differences in geology, relief, drainage, age of land formation and pattern of sedimentary deposition. These differences ultimately influence the nature and properties of the soils in the different physiographic regions.

The biogeochemical cycling of arsenic in soils is strongly affected by other elements. Iron is central due to the strong association between insoluble arsenate and iron(III) oxyhydroxides under aerobic conditions and with the mobilization of iron (II) and arsenite under reducing (that is, paddy) conditions (BGS/DPHE, 2001; Smedley and Kinniburgh, 2002; McArthur et al., 2004; Polizzotto et al., 2005). Manganese oxides also have a similar redox chemistry to iron and are strongly implicated in arsenic immobilization/ mobilization during oxic/ anoxic cycling of paddy sediments (Smedley and Kinniburgh, 2002; Hasan et al., 2007). Arsenate is a phosphate analogue and, thus, key to competition for binding sites within the soil solid phase, as well as having similar biogeochemical cycling under oxic conditions (Adriano, 2001; Meharg and Hartley-Whitaker, 2002; Smith et al., 2002; Lambkin and Alloway, 2003; Stachowicz et al., 2008). Calcium and magnesium immobilize arsenate under oxic conditions, and could also have a role in the biogeochemical cycling of arsenic at a landscape level (Smith et al., 2002; Stachowicz et al., 2008; Fakhreddine et al., 2015).

92

93 Here, we wanted to understand the relationship between soil arsenic and paddy
94 management practice with respect to arsenic loadings in Bangladeshi soils. Cultivation zones
95 of paddy soils (n = 1209) across ten physiographic regions of Bangladesh, from latitude
96 22°06' to 24°53', and longitude 88°20' to 90°59' were sampled and analysed for arsenic and
97 a suite of sixteen other elements (aluminium, calcium, cadmium, cobalt, chromium, copper,
98 iron, lead, magnesium, manganese, molybdenum, nickel, phosphorus, potassium, sodium,
99 and zinc). For a subset of soils (n = 235), paired paddy and adjacent non-paddy soils were
100 also collected and characterised. The data were used to address four specific objectives: to
101 assess the impact that geomorphological differences have on soil arsenic at a landscape
102 level; to understand the relationship between the concentration of arsenic in paddy soils
103 with arsenic in the underlying groundwater; to determine if the source of irrigation water
104 impacts on soil arsenic concentrations; and by examining the concentrations of arsenic and
105 other elements in paddy and non-paddy soils, we aimed to understand the impacts that
106 paddy management has on soil elemental concentrations.

2. Materials and Methods

2.1 Collection of soil samples

A total of 1444 soil samples (topsoil, 0-15 cm from the surface) from paddy fields (n = 1209) and neighbouring non-paddy areas (n = 235) were collected from 10 different physiographic regions within 57 sub-districts (upazilas) from 17 districts of Bangladesh (Table S1). Non-paddy soils were defined as the soils where paddy cultivation and groundwater irrigation had not been practiced within known memory of the farmers. The physiographic regions from where the soil samples were collected included Arial Bil (n = 42 paddy and 10 non-paddy soils), Brahmaputra Floodplain (n = 207 paddy and 64 non-paddy soils), Ganges River Floodplain (n = 261 paddy and 58 non-paddy soils), Ganges Tidal Floodplain (n = 47 paddy and 11 non-paddy soils), Gopalganj-Khulna Bils (n = 63 paddy and 8 non-paddy soils), Karatoya-Bangali floodplain (n = 15 paddy soils only), Meghna Estuarine Floodplain (n = 204 paddy and 28 non-paddy soils), and Meghna River Floodplain (n = 184 paddy and 26 non-paddy soils) from Holocene floodplains, and Barind Tract (n = 68 paddy and 15 non-paddy soils) and Madhupur Tract (n = 118 paddy and 15 non-paddy soils) from Pleistocene terraces. The source of irrigation water for the paddy soils was recorded (groundwater, n = 904; surface water, n = 281; both, n = 24). Only the soils that had a non-mixed irrigation source were used for analyzing the impact of irrigation type on soil arsenic.

2.2 Sample processing and preparation for analysis

The soil samples were air-dried and, prior to analysis, the samples were oven dried (80°C ± 5°C for 48 h), and finely ground using a ball-mill. The soil digestion procedure followed was described by Adomako et al. (2009). Briefly, 0.1 g of soil was placed in a glass digest tube and 2.5 ml of concentrated nitric acid was added to the tube and left overnight for pre-

digestion. Then, 2.5 ml of hydrogen peroxide was added to the sample just before digesting and the sample was heated on the block digester for 1 h at 80°C, for 1 h at 100°C, for 1 h at 120°C, and finally, at 140°C for 3 h until the solution was clear. Once cooled, the digested soil samples were transferred into 15 ml polypropylene tubes and each glass tube was thoroughly rinsed 3 times with ultrapure deionized water (Milli-Q 18.2 MΩ). The volumes were made up to 15 ml mark using the same water. To obtain the appropriate dilution for analysis by inductively coupled plasma-mass spectrometer (ICP-MS) and microwave plasma-atomic emission spectrometer (MP-AES), the samples were further diluted to 1 in 10. Calibration standards were prepared from 1000 mg/l multi-element stock solutions (SPEX CertiPrep Reference Material).

2.3 Chemical analysis

The pH of the soil samples were measured at a soil:water (deionized water) ratio of 1:2.5 (Huq and Alam, 2005). The ICP-MS (Agilent Technologies 7500c, Japan) was used to determine the total concentrations of arsenic, cadmium, cobalt, copper, chromium, lead, manganese, molybdenum, nickel, phosphorus, and zinc in the soil digests and the MP-AES (Agilent Technologies 4100 Series, USA) was used to determine the total concentrations of aluminum, calcium, iron, magnesium, potassium, and sodium in the soil digests. In each batch of digestion, ten percent of the total number of samples were selected randomly for duplicate analysis (n =172). Every batch of samples consisted of 33 randomly selected soil samples, 4 duplicates, 1 blank, and 1 soil CRM (certified reference material) (NCS ZC 73007, China National Analysis Center for Iron and Steel), which were randomized prior to chemical analysis.

155 *2.4 Soil mapping*

156 The data used to perform the mapping of arsenic in paddy soils across Bangladesh included
157 the 1209 paddy soils analyzed in this study as well as 395 soil arsenic concentrations from
158 previous studies (Williams et al., 2011; Lu et al., 2009; Islam et al., 2012). ArcGIS v.10.2 (Esri)
159 was used to create and analyze groundwater and soil arsenic map. The groundwater arsenic
160 data were obtained from BGS/DPHE (2001).

161

162 *2.5 Statistical Analysis*

163 All statistical analyses were performed using the statistical software Minitab v.16 (State
164 College PA) and SigmaPlot v.13 (Systat Software Inc., CA). The data were checked for
165 normality and were transformed prior to statistical analysis where appropriate.

166

3. Results and Discussion

The CRM recovery for arsenic (13.9 ± 0.09 mg/kg, 77.1%, $n = 47$, Table S2) is comparable to other studies using the same methodology (for example, Lu et al., 2010).

To develop a soil arsenic map of the sampled soils, all paddy soil sampling locations within a 10 km² grid were averaged (Fig. 1). Individual locations and sampling densities are shown in Fig. S1. There is a clear north/ south divide in paddy arsenic concentrations with much higher concentrations, in general, in the south. The paddy soil arsenic levels reported here (1-88 mg/kg, average = 8 mg/kg) are within the ranges reported for previous Bangladesh paddy soil surveys (Huq et al., 2003; Meharg and Rahman, 2003; Lu et al., 2009; Williams et al., 2011; Huq and Shoaib, 2013). The pattern of paddy soil concentrations relate well to groundwater measurements (BGS-DPHE, 2001), again with groundwater elevated in the south, excluding the coastal zone. The exception is the cluster of sampling points in the extreme south-east that have a low soil arsenic concentration and the highest groundwater arsenic concentration. This is probably due to the source of irrigation water used in this south-east region, where the main irrigation method is from surface water rather than groundwater (Fig. S2). When comparing the arsenic concentrations in the paddies that have been irrigated with groundwater and surface water across Bangladesh, there was a significant difference ($^{ANOVA} F = 26.23$, $p < 0.001$) in the soil arsenic concentration (Fig. 2). Soils irrigated with groundwater had on average an arsenic concentration of 8.5 mg/kg which was significantly higher than the soils irrigated with surface water, which had an average arsenic concentration of 5.7 mg/kg. As the samples were collected from different geomorphic regions, these results could be confounded by the underlying geomorphology. However, difference in soil arsenic due to different irrigation techniques appears to be a

general trend across the country. For the individual physiographic regions, seven of the regions had groundwater and surface water irrigated soils ($n \geq 10$) to do comparisons between irrigation method and soil arsenic. There was no significant difference in soil arsenic between the groundwater irrigation and surface water irrigation for four of the seven physiographic regions. For the three other physiographic regions, significant differences in arsenic concentrations were observed between the soils irrigated with groundwater (GWI) and surface water (SWI), with higher arsenic concentrations in the groundwater irrigated soils than in the surface water irrigated soils (Ganges Tidal Floodplain, $^{ANOVA}F = 5.97$, $p < 0.05$, $n = 28$ (GWI) and 20 (SWI), mean = 14.6 mg/kg (GWI) and 8.6 mg/kg (SWI); Meghna Estuarine Floodplain, $^{ANOVA}F = 14.84$, $p < 0.001$, $n = 69$ (GWI) and 111 (SWI), mean = 8 mg/kg (GWI) and 3.9 mg/kg (SWI); Meghna River Floodplain, $^{ANOVA}F = 62.06$, $p < 0.001$, $n = 130$ (GWI) and 54 (SWI), mean = 9.4 mg/kg (GWI) and 4.7 mg/kg (SWI)).

Soil arsenic concentrations across ten different physiographic regions of Bangladesh were compared to see how the concentrations varied between the different regions. Significant variations ($^{ANOVA}F = 75.28$, $p < 0.001$ and $^{ANOVA}F = 6.33$, $p < 0.001$, respectively for paddy and non-paddy soils) were observed in soil arsenic concentrations among the ten physiographic regions (Fig. 3 paddy soils; Fig. S3 non-paddy soils). For the paddy soils the Madhupur Tract and the Barind Tract were found to have the lowest mean arsenic concentrations (3.4 mg/kg and 2.8 mg/kg, respectively), whereas the Ganges River Floodplain (11 mg/kg) and the Ganges Tidal Floodplain (13.1 mg/kg) soils had on average the highest soil arsenic concentrations. Martin et al. (2014, 2015) reported higher concentrations and mobilization of arsenic in the Ganges floodplain soils, due to enhanced influence of the pedoenvironmental properties in the region, compared to that in the Meghna floodplain

soils suggesting a complex interaction between soil properties, climate and agricultural management practices in the paddy soil environment in Bangladesh. In the present study, the Ganges floodplain soils were classified as Ganges River Floodplain and Ganges Tidal Floodplain, and the Meghna floodplain soils were classified as Meghna River Floodplain and Meghna Estuarine Floodplain. While no significant difference in arsenic concentrations was observed between Ganges River Floodplain and Meghna River Floodplain soils, there were significance differences between the Ganges Tidal Floodplain and Meghna River Floodplain, and Meghna Estuarine Floodplain soils. Additionally, there was a significant difference between the Ganges River Floodplain and the Meghna Estuarine Floodplain. In all these cases, the soils from the Ganges floodplain had a higher average arsenic concentration than those from the Meghna floodplain (Fig. 3). Similar observations were also reported for groundwater arsenic concentrations across the different geomorphological units of the country (BGS/DPHE, 2001; Ravenscroft, 2001).

At a gross level, high and low groundwater arsenic concentration regions are known to be based on physiographic units, with low concentrations of arsenic in groundwaters in the higher altitude Pleistocene terraces, and at high concentrations in Holocene floodplains (BGS/DPHE, 2001; Smedley and Kinniburgh, 2002; Ahmed et al., 2004; Ravenscroft et al., 2005). The explanation for this is that Pleistocene sediments are more highly weathered and leached of arsenic (Ravenscroft, 2001; Ravenscroft et al., 2005). A recent study on the source of arsenic in the Holocene/ Pleistocene sediments from the Terai plain of Nepal (that stratigraphically resemble Bangladeshi sediments) proposed a number of complex processes which can explain the differences in arsenic concentration between Holocene and Pleistocene sediments (Guillot et al., 2015). However, the river systems of Bangladesh

actively rework the landscape, giving lenses of soil remobilized and re-deposited, interlayering Holocene and Pleistocene soils (BGS/DPHE, 2001; Polizzotto et al., 2005; Meharg et al., 2006; Guillot et al., 2015). It is also known that differential loss of arsenic occurs from groundwater irrigated paddy soils during the subsequent monsoonal floods through partitioning of soil arsenic into overlaying floodwaters (Dittmar et al., 2007; Saha and Ali, 2007; Dittmar et al., 2010; Roberts et al., 2010).

To determine the contribution of both natural soil arsenic concentrations and how paddy management practices have contributed towards the current soil arsenic concentration, paired non-paddy and paddy soils from major physiographic units of Bangladesh were analysed (Fig. 4 and Fig. S4). For the paired analysis of paddy soils and non-paddy soils, all elements were found to be statistically different between the two soil types, except for calcium and phosphorous. All the other elements, except manganese, were higher in the paddy soils compared to the non-paddy soils (Fig S4). There was a significant relationship for soil arsenic between the paddy and non-paddy soils (linear regression $R^2 = 0.26$, $p < 0.001$, $n = 235$) (Table S3). The slope of the overall regression (for all soils) was 1.6:1 for paddy:non-paddy, which indicates a general increase in arsenic of 60 percent in the paddy cultivated soils. The soils from the floodplains and bils (low-lying floodplain) followed a similar pattern as the overall regression regardless if they are from the Brahmaputra, Ganges or Meghna floodplains. Pleistocene terrace soils stand apart and do not follow the overall regression, being both on average lower in arsenic, and having less arsenic accumulation in paddy soils compared to Holocene floodplain soils. A paired t-test of the matching paddy and non-paddy soils for arsenic concentrations within the Pleistocene terrace soils indicated that these soils were significantly different ($p < 0.05$), with the non-paddy soils having elevated

263 arsenic concentrations in comparison to the paddy soils, on an average the non-paddy soils
264 had 19 percent higher arsenic. Pleistocene terrace groundwaters are low in arsenic (Nickson
265 et al., 2000; BGS/DPHE, 2001; Ahmed et al., 2004; Ravenscroft et al., 2005), and thus,
266 irrigation of Pleistocene terrace soils with groundwaters should not lead to elevation in
267 arsenic. As Holocene floodplain groundwaters used in paddy irrigation are elevated in
268 arsenic (Ali et al., 2003; Huq et al., 2003; Meharg and Rahman, 2003; Saha and Ali, 2007; Lu
269 et al., 2009; Huq and Shoaib, 2013), irrigation of paddies with arsenic elevated
270 groundwaters has the potential to lead to build-up in soil arsenic. Arsenic in the non-paddy
271 floodplain soils ranged from 1.8-24.3 mg/kg (mean \pm sd = 5.6 ± 2.9 , coefficient of variation =
272 0.52, n = 205), showing that arsenic is naturally variable in Bangladeshi floodplain soils, with
273 this range being 2-11 mg/kg (mean \pm sd = 3.9 ± 1.8 , coefficient of variation = 0.46, n = 30) for
274 the Pleistocene terrace soils. This emphasises the natural variability in soil arsenic, but that
275 variability is less on Pleistocene terrace soils. It is the Holocene soils/ sediments that are
276 exposed to the active reworking that typifies a dynamic estuarine depositional environment
277 (Sullivan and Aller, 1996; BGS/DPHE, 2001; Polizzotto et al., 2005; Meharg et al., 2006; Lu et
278 al., 2009; Guillot et al., 2015), and this may explain the variability. The inherent differences
279 in the sediments of the floodplain basins deposited from different sources over time,
280 differences in arsenic accumulation/ release equilibria related to the indigenous soil
281 chemistry, residence time, depth and duration of monsoon flood water, rate of particle
282 dispersion, rate of leaching to subsurface, and biovolatilization to the atmosphere can also
283 contribute to explain the variability of arsenic in the floodplain soils of Bangladesh (McLaren
284 et al., 2006; Huq et al. 2008; Khan et al., 2009; Roberts et al., 2010; Mestrot et al., 2011;
285 Brammer, 2012a; Martin et al., 2015). In addition, the diversity and complexity of soils in the
286 floodplains of Bangladesh are influenced by variations in flooding depth within the

inundation land types (Brammer, 1997; Huq et al., 2008). Therefore, the accumulation and release of arsenic in soils vary within the toposequence of a landscape due to variations in relief and soil properties, particularly iron, clay, and organic matter contents (Huq et al., 2008; Brammer, 2012b; Ahmed et al., 2011).

Given that different geomorphic regions within the Holocene floodplain and Pleistocene terrace regions follow the same general trends, with the main differences being between floodplain and terrace, further analysis concentrated on floodplain versus terrace comparisons. For the Pleistocene soils, comparing paddy and non-paddy relationships were seen for all elements tested (Fig. 5). However, it was only for arsenic that paddy soils moved away from a 1:1 relationship, and groundwater is specifically only elevated in arsenic to any significant extent (BGS/DPHE, 2001) with respect to levels already found in soil. This is further evidence, that it is groundwater irrigation *per se*, rather than other aspects of field management, such as fertilizer and manuring practices, that perturb paddy soil arsenic levels compared to non-paddy soils. The depletion in macro-nutrients in Pleistocene sediments, particularly the alkaline earths calcium and magnesium, is most apparent. Arsenic is also positively correlated ($r = 0.3$, $p < 0.001$) with soil pH (Fig. S5), with low pH caused by low calcium and magnesium concentrations, cross confirming the interplay of soils factors correlated with arsenic. Iron and phosphorus, two elements intimately associated with the biogeochemical cycling of arsenic (Fitz and Wenzel, 2002; Smith et al., 2002; Heikens et al., 2007), are also highly depleted in Pleistocene soils. Non-essential aluminium, cadmium, and lead also follow the same trend. It has been demonstrated that pedogenic processes are responsible for the depletion of nutrients within soils over time (Peltzer et al., 2010). Additionally, nutrients can be depleted in soils over shorter periods of

time (Chen et al., 2011). Soils that have been under continuous paddy cropping have been shown to be depleted in key macronutrients in a very short period of time, for example, calcium, magnesium, and sodium have been demonstrated to be rapidly lost in paddy soils within 50 years of rice cultivation (Chen et al., 2011).

The wider characterisation of all the Bangladeshi paddy soils shows that the Pleistocene soils have significantly ($p \leq 0.001$) lower concentrations of all tested elements compared to the Holocene soils (Fig S6). This indicates, again, that Pleistocene soils are less sustainable than Holocene soils with respect to their elemental nutritional qualities. As the soil arsenic concentration is lower in Pleistocene soils it would be expected that the rice plants grown on Pleistocene soils would accumulate less arsenic compared to rice grown on Holocene soils. However, as the Pleistocene soils have lower concentrations of all other elements (Fig. S6), it may be expected that rice grown on these soils would accumulate lower concentrations of nutrients. Rice plants grown on Holocene soils, high in arsenic concentration, are expected to have high arsenic concentrations within the grains. It has been shown that high arsenic concentrations within rice grains impacts negatively on other grain elements (Williams et al., 2009; Norton et al., 2010). Both these situations have the potential to lead to rice grains with lower nutrient concentrations. However, this is not well investigated, and warrants further study in Bangladesh, specifically by wide survey of grain versus soil associations for the primary mineral nutrients of human health importance.

What is apparent from the plots of elemental concentration against arsenic is that Holocene soils have a much wider range of arsenic concentrations at higher concentrations of the other elements compared to Pleistocene soils (Fig 6). That is, there is much greater inherent

variability in arsenic compared to other elements, specifically when other elemental concentrations are high. Groundwater for irrigation is the primary source of arsenic to floodplain paddies that are cropped during the dry season and is well known to elevate arsenic in paddy soils (Ali et al., 2003; Huq et al., 2003; Meharg et al., 2003; Dittmar et al., 2007; Saha and Ali, 2007; Huq, 2008; Lu et al., 2009; Ahmed et al., 2011; Huq and Shoaib, 2013). Paddy soils also have differential interaction with monsoonal floods following dry season application of arsenic, with arsenic capable of partitioning from soils into floodwaters (Dittmar et al., 2007; Saha and Ali, 2007; Dittmar et al., 2010; Roberts et al., 2010). As this interaction between floodwater and soil arsenic will be dependent on soil properties and on the dynamics of floodwater patterns for any specific paddy soil, heterogeneity in arsenic removal is expected. As the paddy soils have a higher arsenic concentration compared to the matched non-paddy soils, it would indicate that this process of loss of arsenic from the soils by monsoonal floods is not sufficient to reduce that arsenic concentration in the paddy soils back to the non-paddy soil background concentration.

When Principle Components Analysis (PCA) is used to look at the interrelationships between arsenic and other elements, the soils cluster into Pleistocene and Holocene using the first and second components (Fig. 7, Table S5). There is some overlap in the middle but this is expected perhaps as the large scales at which physiographic regions are drawn will miss the fine detail on the ground. This is further confounded by the lensing of old soils over new and with the sediment depositional environment also being highly active (Polizzotto et al., 2005; Meharg et al., 2006; Lu et al., 2009; Guillot et al., 2015). The direction of the loadings for the components shows that arsenic trends with most elements, and it is only cadmium and molybdenum that generally differ. The PCA analysis gives further strength to the hypothesis

that arsenic is simply associated with less well weathered/ leached sediments, again suggesting it is either due to the geological newness of Holocene sediments or differences between the sources of sediments that gives rise to the arsenic problems in Bangladesh, and elsewhere (Smedley and Kinniburgh, 2002; McArthur et al., 2004; Nickson et al., 2005; Polya et al., 2005; Berg et al., 2007; Mukherjee et al., 2008; Rowland et al., 2008; Winkel et al., 2008; Guillot et al., 2015).

4. Conclusion

Soil arsenic varied across the physiographic regions of Bangladesh, and overall, the Holocene floodplain soils had a greater arsenic concentration than the Pleistocene terrace soils. Not only do the Holocene floodplain soils have higher concentrations of arsenic, but also they have higher concentrations of other elements. In addition, the overall management of the paddy soils plays a critical role in the loading of arsenic into the soils, with paddy soils having a high concentration of arsenic than non-paddy soils, and soils irrigated with groundwater having a higher concentration of arsenic than those irrigated with surface water. Future studies on how paddy management practices can minimise arsenic accumulation in the soils are essential for mitigating arsenic accumulation in rice grains.

Acknowledgements

This work was done as part of a doctoral fellowship plan funded by the Commonwealth Scholarship Commission in the UK. We gratefully acknowledge the Computer and GIS Unit of Bangladesh Agricultural Research Council (BARC) for providing GIS files for soil mapping. The soil samples were imported into the UK under import license IMP/SOIL/6/2013 issued by

382 Science and Advice for Scottish Agriculture. L.H.E.W. and G.D.J. acknowledge funding by the
383 Swiss National Science Foundation (SNF PP00P2_133619; PP00P2_163747).

384

385 **Appendix A. Supplementary Data**

386 Supplementary data to this article can be found online.

387

388 **References**

389 Adomako, E.E., Solaiman, A.R.M., Williams, P.N., Deacon, C., Rahman, G.K.M.M., Meharg,
390 A.A., 2009. Enhanced transfer of arsenic to grain for Bangladesh grown rice compared to US
391 and EU. *Environ. Int.* 35, 476–479.

392 Adriano, D.C., 2001. *Trace Elements in the Terrestrial Environment*. Springer, New York.

393 Ahmed, K.M., Bhattacharya, P., Hasana, M.A., Akhter, S.H., Alam, S.M.M., Bhuyian, M.A.H.,
394 Imam, M.B., Khan, A.A., Sracek, O., 2004. Arsenic enrichment in groundwater of the alluvial
395 aquifers in Bangladesh: an overview. *Appl. Geochem.* 19, 181–200.

396 Ahmed, Z.U., Panaullah, G.M., DeGloria, S.D., Duxbury, J.M., 2011. Factors affecting paddy
397 soil arsenic concentration in Bangladesh: Prediction and uncertainty of geostatistical risk
398 mapping. *Sci. Tot. Environ.* 412–413, 324–335.

399 Ali, M.A., Badruzzaman, A.B.M., Jalil, M.A., Hossain, M.D., Ahmed, M.F., Masud, A.A.,
400 Kamruzzaman, M., Rahman, M.A., 2003. Arsenic in plant-soil environment in Bangladesh, in:
401 Ahmed, M.F., Ali, M.A., Adeel, Z. (Eds.), *Fate of Arsenic in the Environment*. ITN Centre:
402 Bangladesh University of Engineering and Technology, Dhaka, pp. 85–112.

403 Berg, M., Stengel, C., Trang, P.T.K., Viet, P.H., Sampson, M.L., Leng, M., Samreth, S.,
404 Fredericks, D., 2007. Magnitude of arsenic pollution in the Mekong and Red River Deltas —
405 Cambodia and Vietnam. *Sci. Tot. Environ.* 372, 413–425.

406 BGS/DPHE (British Geological Survey/ Bangladesh Department for Public Health
407 Engineering), 2001. *Arsenic contamination of groundwater in Bangladesh*, Final Report.
408 Kinniburgh, D.G. and Smedley, P.L. (Eds.), British Geological Survey Report WC/00/19. BGS,
409 Keyworth, Vol. 2.

410 Brammer, H., 1996. *The Geography of the Soils of Bangladesh*. The University Press Ltd.,
411 Dhaka.

412 Brammer, H., 2012a. *The Physical Geography of Bangladesh*. The University Press Ltd.,
413 Dhaka.

414 Brammer, H., 2012b. Soil complexity and arsenic mitigation. *Agric. Dev.* 17, 31–35.

415 Chen, L.-M., Zhang, G.-L., Effland, W.R., 2011. Soil characteristic response times and
 416 pedogenic thresholds during the 1000-Year evolution of a paddy soil chronosequence. *Soil*
 417 *Sci. Soc. Am. J.* 75, 1807–1820.

418 Dittmar, J., Voegelin, A., Roberts, L., Hug, S.J., Saha, G.C., Ali, M.A., Badruzzaman, A.B.M.,
 419 Kretzschmar, R., 2007. Spatial distribution and temporal variability of arsenic in irrigated rice
 420 fields in Bangladesh. 2. Paddy soil. *Environ. Sci. Technol.* 41, 5967–5972.

421 Dittmar, J., Voegelin, A., Roberts, L.C., Hug, S.J., Saha, G.C., Ali, M.A., Badruzzaman, A.B.M.,
 422 Kretzschmar, R., 2010. Arsenic accumulation in a paddy field in Bangladesh: Seasonal
 423 dynamics and trends over a three-year monitoring period. *Environ. Sci. Technol.* 44 (8),
 424 2925–2931.

425 FAO/UNDP (Food and Agriculture Organization/ United Nations Development Programme),
 426 1988. Land resources appraisal of Bangladesh for agricultural development, Report 2:
 427 Agroecological Regions of Bangladesh. Report prepared for the Government of the People's
 428 Republic of Bangladesh by the Food and Agriculture Organization of United Nations, United
 429 Nations Development Programme, BGD/81/035, Technical report 2.

430 Fakhreddine, S., Dittmar, J., Phipps, D., Dadakis, J., Fendorf, S., 2015. Geochemical triggers
 431 of arsenic mobilization during managed aquifer recharge. *Environ. Sci. Technol.* 49 (13),
 432 7802–7809.

433 Fitz, W.J., Wenzel, W.W., 2002. Arsenic transformations in the soil-rhizosphere plant
 434 system: fundamentals and potential application to phytoremediation. *J. Biotechnol.* 99,
 435 259–278.

436 Guillot, S., Garçon, M., Weinman, B., Gajurel, A., Tisserand, D., France-Lanord, C., van Geen,
 437 A., Chakraborty, S., Huyghe, P., Upreti, B.N., Charlet, L., 2015. Origin of arsenic in Late
 438 Pleistocene to Holocene sediments in the Nawalparasi district (Terai, Nepal). *Environ. Earth*
 439 *Sci.* 74, 2571–2593.

440 Hasan, M.A., Ahmed, K.M., Sracek, O., Bhattacharya, P., von Brömssen, M., Broms, S.,
 441 Fogelström, J., Mazumder, M.L., Jacks, G., 2007. Arsenic in shallow groundwater of
 442 Bangladesh: investigations from three different physiographic settings. *Hydrogeol. J.* 15,
 443 1507–1522.

444 Heikens, A., Panaullah, G.M., Meharg, A.A., 2007. Arsenic behaviour from groundwater and
 445 soil to crops: Impacts on agriculture and food safety. *Rev. Environ. Contam. Toxicol.* 189,
 446 43–87.

447 Huq, S.M.I., 2008. Fate of arsenic in irrigation water and its potential impacts on food chain,
 448 in: Ahuja, S. (Ed.), *Arsenic Contamination of Groundwater: Mechanism, Analysis, and*
 449 *Remediation*. Wiley, Hoboken, pp. 23–49.

450 Huq, S.M.I., Alam, M.D., 2005. *A Hand Book on Analyses of Soil, Plant, and Water*.
 451 Bangladesh Australia Center for Environmental Research: University of Dhaka, Dhaka,
 452 Bangladesh.

453 Huq, S.M.I., Shoaib, J.U.M., 2013. The Soils of Bangladesh. World soils book series. Springer,
454 Dordrecht.

455 Huq, S.M.I., Rahman, A., Sultana, N., Naidu, R., 2003. Extent and severity of arsenic
456 contamination in soils of Bangladesh, in: Ahmed, M.F., Ali, M.A., Adeel, Z. (Eds.), Fate of
457 Arsenic in the Environment. ITN Centre: Bangladesh University of Engineering and
458 Technology, Dhaka, pp. 69–84.

459 Islam, M.R., Brammer, H., Rahman, G.K.M.M., Raab, A., 2012. Arsenic in rice grown in low-
460 arsenic environments in Bangladesh. Water Qual. Expo. Health 4(4), 197–208.

461 Khan, M.A., Islam, M.R., Panaullah, G.M., Duxbury, J.M., Jahiruddin, M., Loeppert, R.H.,
462 2009. Fate of irrigation-water arsenic in rice soils of Bangladesh. Plant Soil 322 (1), 263–277.

463 Lambkin, D. C., Alloway, B. J., 2003. Arsenate-induced phosphate release from soils and its
464 effect on plant phosphorus. Water Air Soil Pollut. 144, 41–56.

465 Lu, Y., Adomako, E.E., Solaiman, A.R.M., Islam, M.R., Deacon, C., Williams, P.N., Rahman,
466 G.K.M.M., Meharg A.A., 2009. Baseline soil variation is a major factor in arsenic
467 accumulation in Bengal Delta paddy rice. Environ. Sci. Technol. 43, 1724–1729.

468 Lu, Y., Dong, F., Deacon, C., Chen, H.-J., Raab, A., Meharg, A.A., 2010. Arsenic accumulation
469 and phosphorus status in two rice (*Oryza sativa* L.) cultivars surveyed from fields in South
470 China. Environ. Pollut. 158, 1536–1541.

471 Martin, M., Bonifacio, E., Jakeer Hossain, K.M., Imamul Huq, S.M., Barberis, E., 2014. Arsenic
472 fixation and mobilization in the soils of the Ganges and Meghna floodplains. Impact of
473 pedoenvironmental properties. Geoderma 228–229, 132–141.

474 Martin, M., Stanchi, S., Hossain, K.M.J., Imamul Huq, S.M., Barberis, E., 2015. Potential
475 phosphorus and arsenic mobilization from Bangladesh soils by particle dispersion. Sci. Tot.
476 Environ. 53629, 973–980.

477

478 McArthur, J.M., Banerjee, D.M., Hudson-Edwards, K.A., Mishra, R., Purohit, R., Ravenscroft,
479 P., Cronin, A., Howarth, R.J., Chatterjee, A., Talukder, T., Lowry, L., Houghton, S., Chadha,
480 D.K., 2004. Natural organic matter in sedimentary basins and its relation to arsenic in anoxic
481 ground water: the example of West Bengal and its worldwide implications. Appl. Geochem.
482 19, 1255–1293.

483 McLaren, R.G., Megharaj, M., Naidu, R., 2006. Fate of arsenic in the soil environment, in:
484 Naidu, R., Smith, E., Qwens, G., Bhattacharya, P., Nadebaum, P. (Eds.), Managing Arsenic in
485 the Environment: From Soil to Human Health. Collingwood: CSIRO Publishing, pp. 157–182.

486 Meharg, A.A., Hartley-Whitaker, J., 2002. Arsenic uptake and metabolism in arsenic resistant
487 and non-resistant plant species. New Phytol. 154, 29–43.

488 Meharg, A.A., Rahman, M., 2003. Arsenic contamination of Bangladesh paddy field soils:
 489 Implications for rice contribution to arsenic consumption. *Environ. Sci. Technol.* 37 (2),
 490 229–234.

491 Meharg, A.A., Scrimgeour, C., Hossain, S.A., Fuller, K., Cruickshank, K., Williams, P.N.,
 492 Kinniburgh, D.G., 2006. Co-deposition of organic carbon and arsenic in Bengal Delta
 493 aquifers. *Environ. Sci. Technol.* 40, 4928–4935.

494 Meharg, A.A., Zhao, F.J., 2012. *Arsenic & Rice*. Springer, Netherlands.

495 Meharg, A.A., Williams, P.N., Adomako, E., Lawgali, Y.Y., Deacon, C., Villada, A., Cambell,
 496 R.C.J., Sun, G., Zhu, Y.-G., Feldmann, J., Raab, A., Zhao, F.-J., Islam, R., Hossain, S., Yanai, J.,
 497 2009. Geographical variation in total and inorganic arsenic content of polished (white) rice.
 498 *Environ. Sci. Technol.* 43, 1612–1617.

499 Mestrot, A., Feldmann, J., Krupp, E.M., Hossain, M., Roman-Ross, G., Meharg, A.A., 2011.
 500 Field fluxes and speciation of arsines emanating from soils. *Environ. Sci. Technol.* 45,
 501 1798–1804.

502 Mukherjee, A., Bhattacharya, P., Savage, K., Foster, A., Bunschuh, J., 2008. Distribution of
 503 geogenic arsenic in hydrologic systems: controls and challenges. *J. Contam. Hydrol.* 99, 1–7.

504 Nickson, R.T., McArthur, J.M., Ravenscroft, P., Burgess, W.G., Ahmed, K.M., 2000.
 505 Mechanism of arsenic release to groundwater, Bangladesh and West Bengal. *Appl.*
 506 *Geochem.* 15, 403–413.

507 Nickson, R.T., McArthur, J.M., Shrestha, B., Kyaw-Myint, T.O., Lowry, D., 2005. Arsenic and
 508 other drinking water quality issues, Muzaffargarh District, Pakistan. *Appl. Geochem.* 20,
 509 55–68.

510 Norton, G.J., Dasgupta, T., Islam, M.R., Islam, S., Zhao, F.-J., Stroud, J.L., McGrath, S.P.,
 511 Feldmann, J., Price, A.H., Meharg, A.A., 2010. Arsenic influence on cultivar variation in grain
 512 trace-element nutrient content in Bengal Delta grown rice. *Environ. Sci. Technol.* 44,
 513 8284–8288.

514 Peltzer, D.A., Wardle, D.A., Allison, V.J., Baisden, W.T., Bardgett, R.D., Chadwick, O.A.,
 515 Condon, L.M., Parfitt, R.L., Porder, S., Richardson, S.J., Turner, B.L., Vitousek, P.M., Walker,
 516 J., Walker, L.R., 2010. Understanding ecosystem retrogression. *Ecol. Monogr.* 80(4),
 517 509–529.

518 Polizzotto, M.L., Harvey, C.F., Sutton, S.R., Fendorf, S., 2005. Processes conducive to the
 519 release and transport of arsenic into aquifers of Bangladesh. *PNAS* 102, 18819–18823.

520 Polya, D.A., Gault, A.G., Diebe, N., Feldman, P., Rosenboom, J.W., Gilligan, E., Fredericks, D.,
 521 Milton, A.H., Sampson, M., Roland, H.A.L., Lythgoe, P.R., Jones, J.C., Middleton, C., Cooke, D.
 522 A., 2005. Arsenic hazard in shallow Cambodian groundwaters. *Mineral. Mag.* 69, 807–823.

523 Ravenscroft, P., 2001. Distribution of groundwater arsenic in Bangladesh related to geology,
 524 in: Jacks, G., Bhattacharya, P., Khan, A.A. (Eds.), *Groundwater Arsenic Contamination in the*

525 Bengal Delta Plains of Bangladesh. Proceedings of the KTH-Dhaka University Seminar,
 526 University of Dhaka, Bangladesh. KTH Special Publication, TRITA-AMI Report 3084, pp. 41–56.

527 Ravenscroft, P., Burgess, W.G., Ahmed, K.M., Burren, M., Perrin, J., 2005. Arsenic in
 528 groundwater of the Bengal Basin, Bangladesh: Distribution, field relations, and
 529 hydrogeological setting. *Hydrogeol. J.* 13 (5-6), 727–751.

530 Roberts, L.C., Hug, S.J., Dittmar, J., Voegelin, A., Kretzschmar, R., Saha, G.C., Ali, M.A.,
 531 Badruzzaman, A.B.M., 2007. Spatial distribution and temporal variability of arsenic in
 532 irrigated rice fields in Bangladesh. 1. Irrigation water. *Environ. Sci. Technol.* 41, 5960-5966.

533 Roberts, L.C., Hug, S.J., Dittmar, J., Voegelin, A., Kretzschmar, R., Wehrli, B., Cirpka, O.A.,
 534 Saha, G.C., Ali, M.A., Badruzzaman, A.B.M., 2010. Arsenic release from paddy soils during
 535 monsoon flooding. *Nat. Geosci.* 3 (1), 53–59.

536 Rowland, H.A.L., Gault, A.G., Lythgoe, P., Polya, D.A., 2008. Geochemistry of aquifer
 537 sediments and arsenic-rich groundwaters from Kandal Province, Cambodia. *Appl. Geochem.*
 538 23, 3029–3046.

539 Saha, G.C., Ali, M.A., 2007. Dynamics of arsenic in agricultural soils irrigated with arsenic
 540 contaminated groundwater in Bangladesh. *Sci. Tot. Environ.* 379 (2-3), 180–189.

541 Smedley, P.L., Kinniburgh, D.G., 2002. A review of the source, behaviour and distribution of
 542 arsenic in natural waters. *Appl. Geochem.* 17, 517–568.

543 Smith, E., Naidu, R., Alston, A.M., 2002. Chemistry of inorganic arsenic in soils: II. Effect of
 544 phosphorus, sodium, and calcium on arsenic sorption. *J. Environ. Qual.* 31 (2), 557–563.

545 Stachowicz, M., Hiemstra, T., van Riemsdijk, W.H., 2008. Multi-competitive interaction of
 546 As(III) and As(V) oxyanions with Ca^{2+} , Mg^{2+} , PO_4^{3-} , and CO_3^{2-} ions on goethite. *J. Colloid*
 547 *Interface Sci.* 320 (2), 400–414.

548 Sullivan, K.A., Aller, R.C., 1996. Diagenetic cycling of arsenic in Amazon shelf sediments.
 549 *Geochim. Cosmochim. Acta* 60, 1465–1477.

550 Williams, P.N., Islam, S., Islam, R., Jahiruddin, M., Adomako, E., Solieman, A.R.M., Rahman,
 551 G.K.M.M., Lu, Y., Deacon, C., Zhu, Y.-G., Meharg, A.A., 2009. Arsenic limits trace mineral
 552 nutrition (Selenium, Zinc, and Nickel) in Bangladesh rice grain. *Environ. Sci. Technol.* 43,
 553 8430–8436.

554 Williams, P.N., Zhang, H., Davison, W., Meharg, A.A., Hossain, M., Norton, G.J., Brammer, H.,
 555 Islam, M.R., 2011. Organic Matter—Solid Phase Interactions Are Critical for Predicting
 556 Arsenic Release and Plant Uptake in Bangladesh Paddy Soils. *Environ. Sci. Technol.* 45,
 557 6080–6087.

558 Winkel, L., Berg, M., Amini, M., Hug, S.J., Johnson, C.A., 2008. Predicting groundwater
 559 arsenic contamination in Southeast Asia from surface parameters. *Nat. Geosci.* 1, 536–542.





Fig. 1. Sampling locations grouped per 10 km² and sample location marker scaled to size for average arsenic content of that location for surface soils. The underlying contour map is for groundwater arsenic with data inputted from the BGS/DPHE (2001) arsenic survey.



564

565

566

567

568

569

Fig. 2. Box and whisker plot showing concentrations of arsenic in paddy soils irrigated with groundwater and surface water. The boxplots indicate the lower and upper quartile (box), the median (solid line), the mean (dashed line), the 10th and 90th percentiles (whiskers) and the 5th and 95th percentiles (circles). The F-value and p-value from one-way analysis of variance (ANOVA) test are also presented.




Fig. 3: Arsenic concentrations in the paddy soils from different physiographic regions. The numbers of samples (n) at each of the physiographic regions are given within the parentheses. One-way analysis of variance was used to compare pair-wise the means of arsenic concentrations at each of the physiographic regions. Regions that share the same letter (A–E) are not significantly different. The letters indicate Tukey groupings for the physiographic regions with respect to their mean soil arsenic concentrations.





Fig. 4. Relationships between arsenic in paddy and non-paddy soils from different physiographic regions of Bangladesh. The regression line in each graph is the regression line for all the data ($\text{Paddy As} = -0.578 + 1.65 \text{ Non-paddy As}$, $\text{linear regression } R^2 = 0.26$, $p < 0.001$, $n = 235$). The fit and line equations are given in table S3.



583

Non-paddy concentration (mg/kg)

584 **Fig. 5.** Paddy versus non-paddy elemental relationships with soils classified as Holocene and
 585 Pleistocene. The line on each of the graphs is the regression line for each of the elements.
 586 The fit and line equations are given in table S3.




Fig. 6. Relationships for arsenic versus elements for paddy soils grouped into Holocene and Pleistocene. The line on each of the graphs is the regression line for the corresponding elements. The fit and line equations are given in table S4.




Fig. 7. PCA of paddy soils classified into Holocene floodplains and Pleistocene terraces along with loading plot. The first and second component contributed 54.7 and 10.3 percent, respectively, to the variations. The summary of the PCA analysis is given in table S5.